

9. Centrifuge at 7000g at 4°C for 15 min, decant the ethanol, and drain the tube well.
10. Resuspend the pellet in 1 mL H<sub>2</sub>O and add 3 mL of 4M sodium acetate, pH 6.0, and incubate on ice for 2 h. This step precipitates single-stranded RNA but tends to solubilize protein and DNA contaminants that are insoluble in ethanol.
11. Centrifuge as before, drain the pellet well, and resuspend the pellet in 2 mL H<sub>2</sub>O. **Note:** This pellet is more difficult to resuspend than the previous ones, so a larger volume of H<sub>2</sub>O is added. Add 2 vol of ethanol, plus 0.05 vol of 3M sodium acetate, pH 4.8, incubate at -20°C, and centrifuge as before.
12. Resuspend the pellet in 500 µL H<sub>2</sub>O and determine the concentration of RNA by measuring the absorbance at 260 nm and scan the sample from 210–310 nm. 1 OD at 260 nm is equal to a concentration of 40 µg/mL RNA.
13. To check that the RNA is not degraded, analyze a portion on a 1% (w/v) agarose gel prepared in 1X TAE.

#### 4. Notes

1. Hordeivirus particles tend to aggregate because the particles have a strong negative charge and, hence, polycations will interact with them. When performing the virus purification, try to complete it in 1 d, because aggregation of the particles increases as the time used for purification increases.
2. If BSMV is stored in 5% ethylene glycol at -20°C, it remains infectious for many years.

#### Acknowledgments

The financial support for this research was provided by the United States Department of Agriculture from 1978 to the present award, 94-37303-0766.

#### References

1. Hamilton, R. I. and Jackson, A. O. (1995) Hordeivirus, in *Virus Taxonomy, Sixth Report of the International Committee on Taxonomy of Viruses* (Murphy, F. A., Fauquet, C. M., Bishop D. H. L., Ghabrial, S. A., Jarvis, A. W., Martelli, G. P., Mayo, M. A., and Summers, M. D., eds.), Springer-Verlag, Wien New York, pp. 441–444.
2. Gustafson, G. and Armour, S. L. (1986) The complete sequence of RNA  $\beta$  from the type strain of barley stripe mosaic virus. *Nucleic Acids Res.* **14**, 3829–3841.
3. Gustafson, G., Hunter, B., Hanau, R., Armour, S. L., and Jackson, A. O. (1987) Nucleotide sequence and genetic organization of barley stripe mosaic virus RNA  $\gamma$ . *Virology* **158**, 394–406.
4. Gustafson, G., Armour, S. L., Gamboa, G. C., Burgett, S. G., and Shepherd, J. W. (1989) Nucleotide sequence of barley stripe mosaic virus RNA  $\alpha$ : RNA  $\alpha$  encodes a single polypeptide with homology to corresponding proteins from other viruses. *Virology* **170**, 370–377.
5. Petty, I. T. D., French, R., Jones, R. W., and Jackson, A. O. (1990) Identification of barley stripe mosaic virus genes involved in viral RNA replication and movement. *EMBO J.* **9**, 1953–1957.